



**HAL**  
open science

# Repeated maximal-intensity hypoxic exercise superimposed to hypoxic residence boosts skeletal muscle transcriptional responses in elite team-sport athletes

Franck Brocherie, Grégoire Millet, G. d'Hulst, Ruud Van Thienen, Louise Deldicque, Olivier Girard

## ► To cite this version:

Franck Brocherie, Grégoire Millet, G. d'Hulst, Ruud Van Thienen, Louise Deldicque, et al.. Repeated maximal-intensity hypoxic exercise superimposed to hypoxic residence boosts skeletal muscle transcriptional responses in elite team-sport athletes. *Acta Physiologica*, 2018, 222, 10.1111/apha.12851 . hal-02108416

**HAL Id: hal-02108416**

**<https://insep.hal.science//hal-02108416>**

Submitted on 24 Apr 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Received Date : 23-Apr-2016

Revised Date : 30-Nov-2016

Accepted Date : 16-Jan-2017

Article type : Regular Paper

**Repeated maximal-intensity hypoxic exercise superimposed to hypoxic residence boosts skeletal muscle transcriptional responses in elite team-sport athletes**

Franck Brocherie<sup>1, 2</sup>, Grégoire P. Millet<sup>1, 2</sup>, Gommaar D'Hulst<sup>3</sup>, Ruud Van Thienen<sup>3</sup>, Louise Deldicque<sup>3, 4</sup>, Olivier Girard<sup>1, 5</sup>

<sup>1</sup> ISSUL, Institute of Sports Sciences, University of Lausanne, Switzerland,

<sup>2</sup> Department of Physiology, Faculty of Biology and Medicine, University of Lausanne, Switzerland,

<sup>3</sup> Exercise Physiology Research Group, Department of Kinesiology, KU Leuven, Leuven, Belgium,

<sup>4</sup> Institute of Neuroscience, Université catholique de Louvain, Belgium,

<sup>5</sup> ASPETAR, Orthopaedic and Sports Medicine Hospital, Athlete Health and Performance Research Centre, Doha, Qatar.

Running title: Molecular changes to combined hypoxia

Corresponding author:

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi:

10.1111/apha.12851

This article is protected by copyright. All rights reserved.

Prof. Grégoire P. Millet

ISSUL, Building Geopolis, Campus Dorigny, University of Lausanne, CH-1015, Lausanne, Switzerland.

Ph. +41 21 692 32 94

Fax. +41 21 692 32 93

Email gregoire.millet@unil.ch

## ABSTRACT

**Aim.** To determine whether repeated maximal-intensity hypoxic exercise induces larger beneficial adaptations on the hypoxia inducible factor-1 $\alpha$  pathway and its target genes than similar normoxic exercise, when combined with chronic hypoxic exposure.

**Methods.** Lowland elite male team-sport athletes underwent 14 days of passive normobaric hypoxic exposure ( $\geq 14$  h.day<sup>-1</sup> at F<sub>I</sub>O<sub>2</sub> 14.5-14.2%) with the addition of six maximal-intensity exercise sessions either in normobaric hypoxia (F<sub>I</sub>O<sub>2</sub> ~14.2%) (LHTLH; n = 9) or in normoxia (F<sub>I</sub>O<sub>2</sub> 20.9%) (LHTL; n = 11). A group living in normoxia with no additional maximal-intensity exercise (LLTL; n = 10) served as control. Before (Pre), immediately after (Post-1), and 3 weeks after (Post-2) the intervention, muscle biopsies were obtained from the *vastus lateralis*.

**Results.** Hypoxia inducible factor-1 $\alpha$  subunit, vascular endothelial growth factor, myoglobin, peroxisome proliferator-activated receptor-gamma coactivator 1 alpha and mitochondrial transcription factor A mRNA levels increased at Post-1 (all  $P \leq 0.05$ ) in LHTLH, but not in LHTL or LLTL, and returned near baseline levels at Post-2. The protein expression of citrate synthase increased in LHTLH ( $P < 0.001$  and  $P < 0.01$  at Post-1 and Post-2, respectively) and

LLTL ( $P<0.01$  and  $P<0.05$  at Post-1 and Post-2, respectively), whereas it decreased in LHTL at Post-1 and Post-2 (both  $P<0.001$ ).

**Conclusion.** Combined with residence in normobaric hypoxia, repeated maximal-intensity hypoxic exercise induces short-term post-intervention beneficial changes in muscle transcriptional factors that are of larger magnitude (or not observed) than with similar normoxic exercise. The decay of molecular adaptations was relatively fast, with most of benefits already absent 3 weeks post-intervention.

**Key words.** Hypoxia, physical exercise, maximal intensity, gene, protein, oxygen sensor system.

## INTRODUCTION

Human skeletal muscle is a greatly specialized and a highly adaptive tissue. Increased oxygen ( $O_2$ ) consumption and/or a lowered tissue  $O_2$  tension (hypoxia) are known to initiate a cascade of systemic, local and cellular adaptations, all aiming to restore  $O_2$  homeostasis (Semenza, 1999, Semenza, 1998). Whatever the origin of the hypoxic stimulus [*i.e.*, exercise-induced (Ameln *et al.*, 2005) or environmentally- ( $O_2$ -deprived environments) (Hoppeler and Vogt, 2001)], the hypoxia inducible factor-1 $\alpha$  subunit (HIF-1 $\alpha$ ), an  $O_2$  sensitive transcriptional activator that stabilizes in the nucleus under hypoxic conditions, is the main factor mediating these responses (Wang and Semenza, 1995). HIF-1 $\alpha$  is a key regulator responsible for the induction of hypoxia-induced genes (Ke and Costa, 2006) in turn involved in erythropoiesis/iron metabolism, angiogenesis, glucose metabolism as well as cell proliferation/survival and apoptosis (Lundby *et al.*, 2009, Semenza, 1998, Semenza, 1999).

The current scientific literature indicates that prolonged hypoxic residence ( $\geq 10$ -12 h.day<sup>-1</sup> for a minimum of 10 consecutive days) would only induce minimal adaptations in human skeletal muscle tissue. Specifically, down-regulation of HIF-1 $\alpha$  (-49%) and its target genes [*e.g.*, vascular endothelial growth factor (VEGF) (-66%)] (D'Hulst *et al.*, 2016), absence of muscle angiogenesis as well as marginal changes in oxidative enzymes [*e.g.*, citrate synthase (CS)] (Lundby *et al.*, 2009) have been reported. Contrastingly, exercising in hypoxia appears appealing to promote structural and functional adaptations in skeletal muscle (Lundby *et al.*, 2009, Hoppeler and Vogt, 2001). While these changes have been reported to occur immediately after (within days post-intervention) the aforementioned hypoxic interventions, the extent to which delayed (after several weeks) positive muscle phenotype adaptations also occur is unknown.

While chronic low-intensity 'aerobic' exercise in hypoxia may evoke cellular adaptations via HIF-1 $\alpha$  activation (Vogt *et al.*, 2001, Zoll *et al.*, 2006), the magnitude of these responses likely depends on the hypoxic dose (Lundby *et al.*, 2009), and may not necessarily translate in substantial physical performance benefits in endurance individuals (Lundby *et al.*, 2012, Roels *et al.*, 2007, Truijens *et al.*, 2003). Interestingly, it has been postulated that exercise intensity in hypoxia *per se* modulates muscle molecular mechanisms of O<sub>2</sub> homeostasis with '*adaptations that compensate for the reduced availability of O<sub>2</sub> during exercise*' (Hoppeler and Vogt, 2001). Reportedly, maximal-intensity hypoxic exercise, where short 'all-out' efforts ( $\leq 10$  s) with incomplete recoveries ( $< 30$  s) are repeated (namely repeated sprints in hypoxia or RSH), induces additional molecular adaptations at the skeletal muscle level compared to similar exercise in normoxia (RSN) (Faiss *et al.*, 2013b). Specifically, the mRNA expression of genes involved in O<sub>2</sub> signaling (HIF-1 $\alpha$ ), O<sub>2</sub> carrying [myoglobin (Mb)] and pH regulation [carbonic anhydrase-3 (CA-3)] were up-regulated after RSH but not after RSN. Nonetheless, the observation of a concomitant down-regulation of genes involved in

mitochondrial biogenesis [mitochondrial transcription factor A (TFAM) and peroxisome proliferator-activated receptor-gamma coactivator 1 alpha (PGC-1 $\alpha$ )] after RSH would suggest a shift from aerobic to anaerobic glycolytic activity in the muscle and a more efficient use of fast-twitch (FT) muscle fibers (Faiss et al., 2013a, Faiss et al., 2013b, Puype et al., 2013).

We recently demonstrated that the combination of the ‘live high-train low’ (LHTL) paradigm and RSH, namely ‘live high-train low and high’ (LHTLH), produces superior short- (2-3 days post-intervention) and long-term (3 weeks post-intervention) performance benefits (*i.e.*, repeated-sprint ability and high-intensity intermittent exercise capacity) than when LHTL and RSN are combined (Brocherie *et al.*, 2015). While short- and long-term hematological adaptations (*e.g.*, increase in hemoglobin mass) were similar between these two training interventions, twice larger immediate repeated-sprint performance gains associated with LHTLH (which were also maintained after 3 weeks with this training regimen only) advocate that non-hematological factors outside the role played by O<sub>2</sub> carrying-capacity are probably more robust to explain why LHTLH maximizes performance changes. Thus, chronic hypoxic exposure in combination with repeated maximal-intensity hypoxic exercise (LHTLH) appears as a promising intervention to induce concomitant hematological and molecular adaptations. To date, however, changes in skeletal muscle molecular mechanisms of O<sub>2</sub> homeostasis in response to LHTLH remain undetermined.

Therefore, the purpose of this study was to investigate the immediate (*i.e.*, few days post-intervention) and delayed (*i.e.*, several weeks post-intervention) skeletal muscle molecular adaptations associated with 14 days of passive normobaric hypoxic exposure combined with RSH (LHTL+RSH, namely LHTLH) or RSN (LHTL+RSN, namely LHTL) [both compared to a control condition, *i.e.*, ‘live low-train low’ (LLTL)]. Our hypothesis was that, when combined with passive normobaric hypoxic residence, repeated maximal-intensity hypoxic

exercise induces immediate post-intervention muscle molecular adaptations – *i.e.*, HIF-1 $\alpha$  pathway and its target genes – not observed (or to a lower extent) with similar normoxic exercise. We also expected that the molecular adaptations would be maintained for a longer period when combining passive normobaric hypoxic residence with repeated maximal-intensity hypoxic exercise compared to the other conditions tested. This is the first study to investigate the effects of prolonged passive normobaric hypoxic exposure with superimposed repeated maximal-intensity exercise sessions in hypoxia (LHTLH) *vs.* normoxia (LHTL) on molecular adaptations in human skeletal muscle.

## **MATERIALS AND METHODS**

### **Subjects**

Thirty lowland elite male field hockey players (age  $25.1 \pm 4.5$  years, height  $177.8 \pm 5.6$  cm, body weight  $75.2 \pm 7.7$  kg and estimated  $\text{VO}_{2\text{max}}$   $52.0 \pm 1.9$  mL.min<sup>-1</sup>.kg<sup>-1</sup>) were recruited among Belgium, Spanish and Dutch first division clubs to participate in this study.

The subjects were fully informed of the possible risks involved in the study before providing written consent. The study was approved by the Anti-Doping Lab Qatar institutional review board (Agreement SCH-ADL-070) and was conducted according to the Helsinki Declaration.

Exclusion criteria for participation were acclimatization or exposure to hypoxia of more than 2000 m for more than 48 h during a period of 6 months before the study, and any history of altitude-related sickness and health risk that could compromise the subject's safety during exercise and/or hypoxia exposure. During the study, two subjects (control group:  $n = 1$ ; experimental groups:  $n = 1$ ) were excluded due to injury.

## Procedures

The experimental design and physical performance results have been described in details elsewhere (Brocherie *et al.*, 2015). Briefly, it consisted in three testing sessions conducted before (Pre-), 2-3 days (Post-1) and three weeks (Post-2) following the 14-days intervention period. After the completion of Pre-, subjects were randomly assigned to one of the three following groups: 14 days of residence in normobaric hypoxia ( $\geq 14$  h.day<sup>-1</sup> at 2800-3000 m, inspired oxygen fraction (F<sub>i</sub>O<sub>2</sub>) 14.5-14.2%) during which subjects exercised (*i.e.*, regular field hockey practices) at sea level with the addition of six repeated maximal-intensity exercise sessions either in normobaric hypoxia simulating an altitude of 3000 m (LHTL+RSH, namely LHTLH; n = 8) or in normoxia (LHTL+RSN, namely LHTL; n = 11) and control (LLTL; n= 9) where subjects did not performed any additional specific exercise. Importantly, this research was successfully run in a double-blinded, controlled manner (Brocherie *et al.*, 2015).

**Living hypoxic exposure.** The sleeping and recreational hypoxic facilities were fully furnished normobaric hypoxic rooms with O<sub>2</sub>-filtration membrane that reduces the molecular concentration of O<sub>2</sub> in ambient air (CAT system, Colorado Altitude Training, Louisville, Colorado, USA). The two intervention groups (LHTLH and LHTL) were exposed (*i.e.*, from 22:00 to 07:00, from 08:00 to 10:00 and again from 13:00 to 16:00; and were encouraged to spend more time in if desired) to a normobaric hypoxia equivalent to 2500 m (F<sub>i</sub>O<sub>2</sub> 15.1%, BP 768.0 mmHg, P<sub>i</sub>O<sub>2</sub> 108.3 mmHg) for the first 24 h of the intervention period (day 1). Thereafter, the O<sub>2</sub> fraction was further decreased to the equivalent of 2800 m (F<sub>i</sub>O<sub>2</sub> 14.5 ± 0.1%, BP 766.8 ± 1.1 mmHg, P<sub>i</sub>O<sub>2</sub> 104.5 ± 0.6 mmHg; days 2-5) and 3000 m (F<sub>i</sub>O<sub>2</sub> 14.2 ± 0.1%, BP 765.3 ± 1.5 mmHg, P<sub>i</sub>O<sub>2</sub> 101.7 ± 0.8 mmHg; days 6-14). Concentrations of ferritin

( $155.2 \pm 78.7 \mu\text{g}\cdot\text{L}^{-1}$ , range: 45-279  $\mu\text{g}\cdot\text{L}^{-1}$ ) and soluble transferrin receptor ( $256.6 \pm 33.7 \text{mg}\cdot\text{dL}^{-1}$ , range: 202-330  $\text{mg}\cdot\text{dL}^{-1}$ ) measured during the 2-weeks lead-in period at sea level indicated that none of the subjects was iron deficient at the commencement of the study.

**Supervised exercise protocol.** In addition to their usual field hockey practices and matches [carefully monitored and reported in (Brocherie *et al.*, 2015)], subjects of the two intervention groups (LHTLH and LHTL) completed six specific repeated maximal-intensity exercise sessions (with at least 36 h in between) on a synthetic grass ground, inside a 45-m long mobile inflatable simulated hypoxic equipment (Altitude Technology Solutions Pty Ltd, Brisbane, Queensland, Australia), as described elsewhere (Girard *et al.*, 2013). For RSH, ambient air was mixed with nitrogen (from pressurized tanks) to reduce  $F_{iO_2}$  to ~14.2% in order to simulate an altitude of 3000 m. In order to blind subjects to altitude, the system was also run for RSN with normoxic airflow ( $F_{iO_2}$  21.0%) into the tunnel.

Each session lasted ~50 min including a 15-min warm-up, the repeated maximal-intensity exercise and a 10-min recovery phase (*i.e.*, a total of 300 min for the 6 sessions among the 14-days intervention). Specifically, the repeated maximal-intensity exercise included 4 sets of 5 × 5-s maximal sprints interspersed with 25 s of passive recovery with 5 min of standing rest between sets.

### **Muscle Biopsy Samples**

Biopsy samples were all taken by the same experienced medical doctor with a 5-mm Bergström type needle (in conjunction with a suction device to create a negative pressure) in the mid portion of the *vastus lateralis* muscle after local anesthesia (1% xylocaine,

subcutaneously). Biopsies were preceded by 48 h without any exercise activity and were taken randomly in opposite legs (*i.e.*, left-right-left or vice versa) during subsequent test sessions. For mRNA analysis, the muscle tissue portion was immediately frozen in liquid nitrogen and stored at -80°C until required for analyses.

### **Western blotting**

Frozen muscle tissue (~20 mg) was homogenized 3 × 5 s with a TissueLyser (Qiagen, Hilden, Germany) in an ice-cold buffer (1:10, w/v) [50 mM Tris-HCl pH 7.0, 270 mM sucrose, 5 mM EGTA, 1 mM EDTA, 1 mM sodium orthovanadate, 50 mM glycerophosphate, 5 mM sodium pyrophosphate, 50 mM sodium fluoride, 1 mM DTT, 1 % Triton-X 100 and a protease inhibitor cocktail (Roche Applied Science, Vilvoorde, Belgium)]. After centrifugation of homogenates at 10 000 g for 10 min at 4°C, the supernatants were stored at -80°C. Protein concentration was measured using the DC protein assay kit (Bio-Rad laboratories, Nazareth, Belgium) with bovine serum albumin as a standard. Proteins (30-50 µg) were separated by SDS-PAGE (7.5 – 12.5%) and transferred to PVDF membranes. Subsequently, membranes were blocked with 5% non-fat milk for 1 h and incubated overnight (4°C) with the following antibodies: glucose transporter 4 (GLUT-4, #PA1-1065, Thermo Scientific, Erembodegem, Belgium), phosphofructo kinase (PFK, #166722, Santa Cruz, Huissen, The Netherlands), monocarboxylate transporter-1 (MCT-1, #AB3538P, Millipore, Overijse, Belgium), monocarboxylate transporter-4 (MCT-4, #AB3316P, Millipore), CA-3 (#135995, Abcam, Cambridge, UK), AMP-activated protein kinase alpha (AMPK $\alpha$ , #2532, Cell Signaling, Leiden, The Netherlands), phospho-AMPK $\alpha$  Thr172 (#2535, Cell Signaling), CS (#14309, Cell Signaling), eukaryotic elongation factor 2 (eEF2, #2332, Cell Signaling). Appropriate horseradish peroxidase-conjugated secondary antibodies (Sigma-Aldrich, Bornem, Belgium)

were used for chemiluminescent detection of the proteins of interest. Membranes were scanned and quantified with Genesnap and Genetools softwares (Syngene, Cambridge, UK), respectively. The results are presented as the ratio protein of interest/eEF2 or as the ratio phosphorylated/total form for AMPK.

### **Real-time quantitative Polymerase Chain Reaction**

RNA was extracted using TRIzol (Invitrogen, Vilvoorde, Belgium) from 20–25 mg of frozen muscle tissue. RNA quality and quantity were assessed by spectrophotometry with a Nanodrop (Thermo Scientific, Erembodegem, Belgium). One  $\mu\text{g}$  of RNA was reverse-transcribed using the High-Capacity cDNA Reverse Transcription kit (Applied Biosystems, Gent, Belgium) according to manufacturer's instructions. A SybrGreen-based master mix (Applied Biosystems, Erembodegem, Belgium) was used for real-time PCR analyses using the ABIPRISM 7300 (Applied Biosystems). Real-time PCR primers were designed for human HIF-1 $\alpha$ , PHD-2 (prolyl hydroxylase domain protein 2), VEGF, Mb, cytochrome c oxidase subunit 4 isoform 1 (COX-4\_1) and isoform 2 (COX-4\_2), PGC-1 $\alpha$ , TFAM, endothelial nitric oxide synthase (eNOS), neuronal nitric oxide synthase (nNOS), pyruvate dehydrogenase kinase 1 (PDK-1), glucose transporter 1 (GLUT-1) and, phosphofructokinase (PFK) (Table 1). Thermal cycling conditions consisted of 40 three-step cycles including denaturation of 30 s at 95°C, annealing of 30 s at 58°C and extension of 30 s at 72°C. All reactions were performed in triplicate. To compensate for variations in input RNA amounts and efficiency of reverse transcription ribosomal protein L19 (RPL19) and beta-2-microglobulin (B2M) mRNA were quantified, and results were normalized to these values. These genes were chosen out of three normalization genes using the GeNorm applet according to the guidelines and theoretical framework described elsewhere (Vandesompele *et al.*, 2002).

### **Citrate synthase activity**

Five  $\mu\text{g}$  of protein were added to the reaction buffer containing 100 mM Tris base, 2 mM EDTA, 1.25 mM malate, 0.25 mM NAD and 6  $\text{U}\cdot\text{ml}^{-1}$  L-malate dehydrogenase (10127256001, Roche, Brussels, Belgium). The temperature was set at 37°C and the reaction started by the addition of acetyl-CoA. NADH production was measured by fluorescence (340 nm excitation, 460 nm emission).

### **Data and statistical analysis**

Because of large inter-individual differences in absolute mRNA levels and protein expressions, post-intervention values are expressed as percent changes in means (Pre- vs. Post-1 and Post-2, respectively), while the mean of the Pre- values was assigned to the arbitrarily value of  $1.00\pm 0.00$ . Two-way ANOVA with repeated measures [Time (Pre- vs. Post-1 vs. Post-2)  $\times$  Group (LHTLH vs. LHTL vs. LLTL)] was used to compare each measured variable. When significant modifications were found, Holm-Sidak post-hoc test was performed to localize the effect. All analyses were made using Sigmaplot 11.0 software (Systat Software, CA, USA). Null hypothesis was rejected at  $P < 0.05$ .

## **RESULTS**

### **Skeletal muscle mRNA expression analysis and enzyme activity**

Figures 1, 2 and 3 and Table 2 display changes from Pre- to Post-1 and Post-2 in mRNA and protein expression/activity levels in the three groups.

*O<sub>2</sub> signaling.* Compared to Pre-, HIF-1 $\alpha$  mRNA levels increased at Post-1 ( $P < 0.01$ ) in LHTLH only, before returning to near Pre- values at Post-2 (Fig. 1). At Post-1, higher HIF-1 $\alpha$  mRNA levels were observed in LHTLH compared to LHTL and LLTL (both  $P < 0.05$ ).

*O<sub>2</sub> carrier.* Compared to Pre-, higher mRNA levels of VEGF ( $P = 0.05$ ) and Mb ( $P < 0.05$ ) occurred at Post-1 in LHTLH (Fig. 1). At Post-1, VEGF and Mb values were also higher in LHTLH than in LLTL (both  $P < 0.05$ ).

*Mitochondrial biogenesis and metabolism.* Transcript levels of regulators of mitochondrial biogenesis PGC-1 $\alpha$  ( $P < 0.05$  at Post-1) and TFAM ( $P < 0.05$  at Post-1 and Post-2) were increased in LHTLH only. No significant changes were detected in COX-4\_1 and COX-4\_2 mRNA levels. The protein expression of CS increased in LHTLH ( $P < 0.001$  at Post-1 and  $P < 0.01$  at Post-2) and LLTL ( $P < 0.01$  at Post-1, and  $P < 0.05$  at Post-2), whereas it decreased in LHTL ( $P < 0.001$  at Post-1 and Post-2) (Fig. 2). Similar results were obtained for CS activity (Table 2).

*Nitric oxide synthase pathway.* Compared to Pre-, eNOS and nNOS mRNA levels tended to increase at Post-1 in LHTLH ( $P = 0.30$  and  $P = 0.09$ , respectively) (Fig. 1). In LHTL, only eNOS mRNA levels increased significantly from Pre- to Post-1 ( $P < 0.01$ ), while values of both eNOS and nNOS were higher at Post-2 vs. Pre (both  $P < 0.05$ ). The mRNA levels of eNOS at Post-1 were higher in LHTLH and LHTL compared to LLTL (both  $P < 0.05$ ).

*pH regulation.* The protein expression of MCT-1, but not of MCT-4 and CA-3, was increased in LHTLH ( $P < 0.05$ ) and LHTL ( $P < 0.05$ ) at Post-1 in reference to Pre- (Figs. 2 and 3). Whereas MCT-1 at Post-2 returned to near Pre- values in LHTLH, lower values were observed from Pre- to Post-2 in LHTL ( $P < 0.05$ ).

*Glucose metabolism.* At Post-1, the mRNA levels of GLUT-1 were higher in LHTLH compared to Pre- (Fig. 1,  $P < 0.05$ ) while the protein expression of PFK was lower in LHTLH

compared to LLTL (Figs. 2 and 3,  $P < 0.05$ ). No changes were measured in the protein expression levels of GLUT-4 (Figs. 2 and 3).

*AMPK total protein content and phosphorylation status.* Compared to Pre-, AMPK total protein content ( $P < 0.01$  for LHTLH and LHTL) and pAMPK-tot ( $P < 0.001$  for LHTLH and LHTL) increased significantly at Post-1, whereas lower pAMPK-tot values were noted at Post-2 ( $P < 0.01$  and  $< 0.001$  for LHTLH and LHTL, respectively) (Figs. 2 and 3).

## DISCUSSION

This is the first study to investigate the effects of prolonged passive exposure to normobaric hypoxia with superimposed repeated maximal-intensity exercise sessions in hypoxia (LHTLH) vs. normoxia (LHTL) on molecular regulations in human skeletal muscle. The novel findings are that LHTLH elicits higher short-term (first few days post-intervention) molecular responses of factors implicated in the regulation of O<sub>2</sub> signaling and carrying, mitochondrial biogenesis, as well as of enzymes implicated in mitochondrial metabolism compared to LHTL [with also no change in control (LLTL)]. We further indicate that the majority of these positive molecular responses disappeared already three weeks post-intervention. We confirm our hypothesis of larger specific transcriptional responses when passive normobaric hypoxic exposure and repeated maximal-intensity hypoxic exercise are combined, yet with normalization of molecular adaptations three weeks after the intervention. These adaptations in O<sub>2</sub> signaling and transport, mitochondrial biogenesis, as well as in enzymes implicated in mitochondrial metabolism, may contribute to the sport-specific performance gains (Post-1) reported elsewhere (Brocherie et al., 2015). The functional benefits (*e.g.*, performance gains) were maintained up to 3 weeks post-LHTLH (Brocherie et

al., 2015) and were likely caused by the molecular adaptations which may have promoted changes in muscle biochemistry and vascularity.

### **O<sub>2</sub> signaling and gene regulation**

In the present study, the level of HIF-1 $\alpha$  mRNA increased immediately after the intervention in LHTLH but, unexpectedly, it was not the case for LHTL. First, the absence of HIF-1 $\alpha$  increase after 14 days of LHTL supports the hypothesis that HIF-1 $\alpha$  response to hypoxia is time-dependent and activated shortly and briefly after hypoxic stimulus exposure (Lundby *et al.*, 2009). Accordingly, it has been previously demonstrated that HIF-1 $\alpha$  protein levels peak within the first hours of hypoxic exposure then progressively decline toward basal levels (Stroka *et al.*, 2001, Vigano *et al.*, 2008), suggesting a possible local or systemic ‘acclimatization’ after several days (Lundby *et al.*, 2009). As the protein level of HIF-1 $\alpha$  is only briefly increased after a hypoxic stimulus and our intention was to determine stable molecular changes after repeated expositions to hypoxia, we solely quantified the mRNA level of this transcription factor, which reflects the long-term activation of the HIF-1 $\alpha$  pathway (Galban and Gorospe, 2009). With this in mind, the enhanced mRNA level of HIF-1 $\alpha$  after LHTLH clearly suggests that the addition of repeated maximal-intensity exercise in hypoxia (but not in normoxia) plays an important role for up-regulating the activation of the HIF-1 $\alpha$  pathway and some of its downstream genes (Zoll *et al.*, 2006, Vogt *et al.*, 2001, Faiss *et al.*, 2013b). This is further supported by the return to near HIF-1 $\alpha$  mRNA Pre- values at 3 weeks post-intervention when no additional repeated maximal-intensity exercises and/or hypoxic stimulation were performed.

Activation of HIF-1 $\alpha$  is known to lead to cellular adaptations [*i.e.*, O<sub>2</sub> carrying-capacity (Wenger and Gassmann, 1997), neovascularization (Forsythe *et al.*, 1996), glucose oxidation

(Wenger and Gassmann, 1997)], which in turn would positively influence exercise capacity in humans (Vogt et al., 2001, Zoll et al., 2006, Faiss et al., 2013b). Accordingly, the mRNA levels for the capillary growth factor VEGF, *i.e.*, an HIF-1-regulated gene (Semenza, 1999, Semenza et al., 1999, Wenger and Gassmann, 1997) and Mb mRNA, significantly increased after LHTLH, whereas no changes were observed in both LHTL and LLTL. This corroborates previous works, which demonstrated Mb mRNA or protein levels enhancement after hypoxic endurance exercise [*i.e.*, 3-5 sessions of intermittent hypobaric (~2300 m) or normobaric hypoxic (~3850 m) training per week for 4-6 weeks (Vogt et al., 2001, Terrados et al., 1990)]. Overall, it seems that these RSH-related molecular adaptations were not blunted by the passive normobaric residence and may participate in the twice-larger repeated-sprint performance improvements observed in LHTLH vs. LHTL (Brocherie et al., 2015).

### **Metabolic Phenotype**

After hypoxic endurance exercise, it has been demonstrated that mRNA levels for PGC-1 $\alpha$  (Zoll *et al.*, 2006), COX-1, COX-4 and CS (Zoll et al., 2006, Vogt et al., 2001, Terrados et al., 1990) increase to a greater extent compared to similar intervention in normoxia. In the present study, LHTLH induced larger mitochondrial adaptations (*i.e.*, increased mRNA levels for PGC-1 $\alpha$  and TFAM) compared to LHTL and LLTL, thereby suggesting a preponderant role of the superimposed RSH for muscle phenotypic adaptations. Whereas the protein expression and protein activity of CS decreased after LHTL, it increased after both LHTLH and LLTL. Conversely, when used in isolation (*i.e.*, with normoxic residence), Faiss et al. (Faiss *et al.*, 2013b) indicated that RSH induced a down-regulation in mitochondrial biogenesis (PGC-1 $\alpha$  and TFAM), despite unchanged oxidative capacity (CS). Some methodological differences (*i.e.*, exercise mode, frequency and duration) between studies might be responsible for these

discrepant findings. In summary, the increase in PGC-1 $\alpha$  and TFAM measured after intervention in LHTLH, for which mRNA expression predicts protein levels quite well (Collu-Marchese et al., 2015, Handschin et al., 2003), appears also as a plausible molecular explanation for the beneficial adaptations observed in LHTLH vs. LHTL vs. LLTL.

### **Compensatory vasodilation**

Among the HIF-1 $\alpha$  target genes, VEGF is a critical signal in vascular remodeling, which maintains vascular integrity and stimulates the production of the vasodilatory mediator nitric oxide. Hence, eNOS plays a key role in blood flow regulation and vascular tone (Gielen *et al.*, 2011). Concomitantly, nNOS expression exerts a functionally significant effect in hypoxic tissue, thereby influencing tissue O<sub>2</sub> delivery (Fish et al., 2007, Tsui et al., 2014), ventilatory regulation and metabolic adaptations to hypoxia (Gardiner *et al.*, 2011). Although we did not measure the protein expression, an increase in eNOS and nNOS may promote changes in the blood flow and vascular tone (Gielen *et al.*, 2011), participate in angiogenesis (Viboolvorakul and Patumraj, 2014) and enhance O<sub>2</sub> delivery.

### **pH-regulating System**

In line with previous sprint interval intervention studies (Puype et al., 2013, Burgomaster et al., 2007), LHTLH and LHTL increased muscle MCT-1 protein content, whereas MCT-4 and CA-3 contents did not change. This potential up-regulation of lactate metabolism may reflect the similar sport-specific aerobic performance gains observed in LHTLH and LHTL (Brocherie et al., 2015).

## Delayed effects on skeletal muscle transcriptional regulation

We recently demonstrated that LHTLH and LHTL resulted in similar short-term (2-3 days post-intervention) hematological adaptations (*i.e.*, increase in hemoglobin mass) that were maintained for at least 3 weeks post-intervention (Brocherie *et al.*, 2015). Contrastingly, our novel findings indicate that most of the transcriptional adaptations were back to near Pre-values 3 weeks post-intervention. The present study is the first to show the time course of HIF-1 $\alpha$  and related genes transcription in human skeletal muscle adaptations following residence in normobaric hypoxia superimposed with repeated maximal-intensity exercise in hypoxia or normoxia. With the post-intervention carefully supervised and controlled (*i.e.*, similar between groups), the fact that this period did not include any environmental hypoxic stress – be it during residence or exercise in normobaric hypoxia – indicates that the lack of external hypoxic ‘stimulus’ could be responsible for the rapid reversal of skeletal muscle transcriptional regulation. Meanwhile, it may have already promoted some beneficial muscle function adjustments and therefore functional benefits without affecting hemoglobin mass improvement (Brocherie *et al.*, 2015). Additionally, as the post-intervention period (3 weeks) did not include any repeated maximal-intensity exercise session (as performed six times during the 14-days intervention), the influence of such exercise modality and/or frequency on normalization of molecular responses should not be overlooked. One aspect that deserves more research attention is whether additional hypoxic stress and/or intense exercise when individuals return to sea level after an hypoxic intervention induces a better maintenance of molecular adaptations. The decay of molecular responses over time after return to sea level, that occur more rapidly than the hematological adaptations, would suggest that mechanisms (not measured here) such as those involved in the factor-inhibiting HIF regulation (Lindholm and Rundqvist, 2016) may be at play.

## Strengths and limitations

One potential strength of this study is that elite field hockey players volunteered to participate, while mechanistic studies are generally carried out with recreational participants. In these studies, results are often extrapolated to elite athletes while exercise-related adaptations are probably less prominent as the level of practice/performance increases. Here, we found that LHTLH was the most efficient strategy to activate in the short-term the transcription of specific genes involved in key physiological processes for adaptations such as angiogenesis or O<sub>2</sub> transport. It is therefore tempting to associate the molecular regulations we measured at Post-1 and the enhanced performance that we previously reported at both Post-1 and Post-2 in LHTLH (Brocherie *et al.*, 2015).

Methodologically, it is known that most gene expressions generally peak within 2-8 h after a single exercise stimulus (Pilegaard *et al.*, 2003). Although we acknowledge that short-term (hours) molecular responses are the basis for long term structural, enzymatic and functional adaptations, we are confident that, with muscle biopsies obtained 48 h following the last exercise session in our subjects, the augmented expression of several genes actually is the result of the proposed intervention *per se*, and not a side effect of the final exercise session.

Despite careful control of biopsy sampling (*i.e.*, no physical activity allowed between the last exercise session, similar sampling timing between 08:00 and 10:00 a.m. during the three test sessions and corresponding samples ran in the same assay) (Fluck *et al.*, 2005), a large inter-subject gene and protein expression variation occurred here. Potentially, this may be explained by: (i) DNA sequence variations resulting from HIF-1 $\alpha$  gene polymorphism in the promoter region of HIF-1 $\alpha$  gene (Prior *et al.*, 2006), (ii) fiber type differences in the sampling from *vastus lateralis* muscle, which has an unequal typology and metabolic properties (Pette, 1985), (iii) pulsative nature of gene expression in muscular fiber (Newlands *et al.*, 1998), and (iv) involvement of reactive oxygen species in the regulation of HIF-1 $\alpha$  mRNA and HIF-1 $\alpha$

target genes (Pialoux *et al.*, 2009). That said, it is worth mentioning that biopsies obtained alternatively from the left and right *vastus lateralis* muscle actually give similar mRNA expression profiles when the two legs are compared (Lundby *et al.*, 2005).

In conclusion, combined with normobaric hypoxic residence, repeated maximal-intensity hypoxic exercise elicited higher short-term (first few days post-intervention) skeletal muscle molecular beneficial adaptations not observed (or with smaller magnitudes) after similar normoxic exercise. The large and specific adaptations in mRNA levels of factors involved in O<sub>2</sub> signaling and transport, mitochondrial biogenesis, as well as in enzymes implicated in mitochondrial metabolism, are seen to highlight the prominence of superimposed repeated maximal-intensity hypoxic exercise, yet with a rapid decay and normalization of molecular adaptations after cessation of the intervention.

### **Acknowledgements**

This research was funded by a Grant awarded by Aspetar (Qatar Orthopedic and Sports Medicine Hospital) at the Aspire Zone Foundation, Qatar (AF/C/ASP1905/11). The authors thank Mr. Julien Rysman for his support in the organization of the study and assistance during the experiments and Mr. Damien Naslain for technical assistance for the biochemical assays. The authors declare no conflict of interest, or financial ties to disclose.

### **Conflict of interest**

No conflict of interest, financial or otherwise, is declared by the authors.

## REFERENCES

- Ameln, H., Gustafsson, T., Sundberg, C. J., Okamoto, K., Jansson, E., Poellinger, L. & Makino, Y. 2005. Physiological activation of hypoxia inducible factor-1 in human skeletal muscle. *FASEB J*, 19, 1009-11.
- Brocherie, F., Millet, G. P., Hauser, A., Steiner, T., Rysman, J., Wehrlin, J. P. & Girard, O. 2015. "Live High-Train Low and High" Hypoxic Training Improves Team-Sport Performance. *Med Sci Sports Exerc*, 47, 2140-9.
- Burgomaster, K. A., Cermak, N. M., Phillips, S. M., Benton, C. R., Bonen, A. & Gibala, M. J. 2007. Divergent response of metabolite transport proteins in human skeletal muscle after sprint interval training and detraining. *Am J Physiol Regul Integr Comp Physiol*, 292, R1970-6.
- Collu-Marchese, M., Shuen, M., Pauly, M., Saleem, A. & Hood, D. A. 2015. The regulation of mitochondrial transcription factor A (Tfam) expression during skeletal muscle cell differentiation. *Biosci Rep*, 35.
- D'Hulst, G., Ferri, A., Naslain, D., Bertrand, L., Horman, S., Francaux, M., Bishop, D. J. & Deldicque, L. 2016. Fifteen days of 3,200 m simulated hypoxia marginally regulates markers for protein synthesis and degradation in human skeletal muscle. *Hypoxia*, 4, 1-14.
- Faiss, R., Girard, O. & Millet, G. P. 2013a. Advancing hypoxic training in team sports: from intermittent hypoxic training to repeated sprint training in hypoxia. *Br J Sports Med*, 47 Suppl 1, i45-i50.
- Faiss, R., Leger, B., Vesin, J. M., Fournier, P. E., Eggel, Y., Deriaz, O. & Millet, G. P. 2013b. Significant molecular and systemic adaptations after repeated sprint training in hypoxia. *PLoS ONE*, 8, e56522.
- Fish, J. E., Matouk, C. C., Yeboah, E., Bevan, S. C., Khan, M., Patil, K., Ohh, M. & Marsden, P. A. 2007. Hypoxia-inducible expression of a natural cis-antisense transcript inhibits endothelial nitric-oxide synthase. *J Biol Chem*, 282, 15652-66.
- Fluck, M., Dapp, C., Schmutz, S., Wit, E. & Hoppeler, H. 2005. Transcriptional profiling of tissue plasticity: role of shifts in gene expression and technical limitations. *J Appl Physiol (1985)*, 99, 397-413.
- Forsythe, J. A., Jiang, B. H., Iyer, N. V., Agani, F., Leung, S. W., Koos, R. D. & Semenza, G. L. 1996. Activation of vascular endothelial growth factor gene transcription by hypoxia-inducible factor 1. *Mol Cell Biol*, 16, 4604-13.
- Galban, S. & Gorospe, M. 2009. Factors interacting with HIF-1alpha mRNA: novel therapeutic targets. *Curr Pharm Des*, 15, 3853-60.
- Gardiner, B. S., Smith, D. W., O'Connor, P. M. & Evans, R. G. 2011. A mathematical model of diffusional shunting of oxygen from arteries to veins in the kidney. *Am J Physiol Renal Physiol*, 300, F1339-52.
- Gielen, S., Sandri, M., Erbs, S. & Adams, V. 2011. Exercise-induced modulation of endothelial nitric oxide production. *Curr Pharm Biotechnol*, 12, 1375-84.
- Girard, O., Brocherie, F. & Millet, G. P. 2013. On the use of mobile inflatable hypoxic marquees for sport-specific altitude training in team sports. *Br J Sports Med*, 47 Suppl 1, i121-i123.
- Handschin, C., Rhee, J., Lin, J., Tarr, P. T. & Spiegelman, B. M. 2003. An autoregulatory loop controls peroxisome proliferator-activated receptor gamma coactivator 1alpha expression in muscle. *Proc Natl Acad Sci U S A*, 100, 7111-6.
- Hoppeler, H. & Vogt, M. 2001. Muscle tissue adaptations to hypoxia. *J Exp Biol*, 204, 3133-9.
- Ke, Q. & Costa, M. 2006. Hypoxia-inducible factor-1 (HIF-1). *Mol Pharmacol*, 70, 1469-80.

- Lindholm, M. E. & Rundqvist, H. 2016. Skeletal muscle hypoxia-inducible factor-1 and exercise. *Exp Physiol*, 101, 28-32.
- Lundby, C., Calbet, J. A. & Robach, P. 2009. The response of human skeletal muscle tissue to hypoxia. *Cell Mol Life Sci*, 66, 3615-23.
- Lundby, C., Millet, G. P., Calbet, J. A., Bartsch, P. & Subudhi, A. W. 2012. Does 'altitude training' increase exercise performance in elite athletes? *Br J Sports Med*, 46, 792-5.
- Lundby, C., Nordsborg, N., Kusuhara, K., Kristensen, K. M., Neuffer, P. D. & Pilegaard, H. 2005. Gene expression in human skeletal muscle: alternative normalization method and effect of repeated biopsies. *Eur J Appl Physiol*, 95, 351-60.
- Newlands, S., Levitt, L. K., Robinson, C. S., Karpf, A. B., Hodgson, V. R., Wade, R. P. & Hardeman, E. C. 1998. Transcription occurs in pulses in muscle fibers. *Genes Dev*, 12, 2748-58.
- Pette, D. 1985. Metabolic heterogeneity of muscle fibres. *J Exp Biol*, 115, 179-89.
- Pialoux, V., Mounier, R., Brown, A. D., Steinback, C. D., Rawling, J. M. & Poulin, M. J. 2009. Relationship between oxidative stress and HIF-1 alpha mRNA during sustained hypoxia in humans. *Free Radic Biol Med*, 46, 321-6.
- Pilegaard, H., Saltin, B. & Neuffer, P. D. 2003. Exercise induces transient transcriptional activation of the PGC-1alpha gene in human skeletal muscle. *J Physiol*, 546, 851-8.
- Prior, S. J., Hagberg, J. M., Paton, C. M., Douglass, L. W., Brown, M. D., McLenithan, J. C. & Roth, S. M. 2006. DNA sequence variation in the promoter region of the VEGF gene impacts VEGF gene expression and maximal oxygen consumption. *Am J Physiol Heart Circ Physiol*, 290, H1848-55.
- Puype, J., Van Proeyen, K., Raymackers, J. M., Deldicque, L. & Hespel, P. 2013. Sprint Interval Training in Hypoxia Stimulates Glycolytic Enzyme Activity. *Med Sci Sports Exerc*, 45, 2166-74.
- Roels, B., Bentley, D. J., Coste, O., Mercier, J. & Millet, G. P. 2007. Effects of intermittent hypoxic training on cycling performance in well-trained athletes. *Eur J Appl Physiol*, 101, 359-68.
- Semenza, G. L. 1998. Hypoxia-inducible factor 1: master regulator of O<sub>2</sub> homeostasis. *Curr Opin Genet Dev*, 8, 588-94.
- Semenza, G. L. 1999. Regulation of mammalian O<sub>2</sub> homeostasis by hypoxia-inducible factor 1. *Annu Rev Cell Dev Biol*, 15, 551-78.
- Semenza, G. L., Agani, F., Iyer, N., Kotch, L., Laughner, E., Leung, S. & Yu, A. 1999. Regulation of cardiovascular development and physiology by hypoxia-inducible factor 1. *Ann N Y Acad Sci*, 874, 262-8.
- Stroka, D. M., Burkhardt, T., Desbaillets, I., Wenger, R. H., Neil, D. A., Bauer, C., Gassmann, M. & Candinas, D. 2001. HIF-1 is expressed in normoxic tissue and displays an organ-specific regulation under systemic hypoxia. *FASEB J*, 15, 2445-53.
- Terrados, N., Jansson, E., Sylven, C. & Kaijser, L. 1990. Is hypoxia a stimulus for synthesis of oxidative enzymes and myoglobin? *J Appl Physiol (1985)*, 68, 2369-72.
- Truijens, M. J., Toussaint, H. M., Dow, J. & Levine, B. D. 2003. Effect of high-intensity hypoxic training on sea-level swimming performances. *J Appl Physiol*, 94, 733-43.
- Tsui, A. K., Marsden, P. A., Mazer, C. D., Sled, J. G., Lee, K. M., Henkelman, R. M., Cahill, L. S., Zhou, Y. Q., Chan, N., Liu, E. & Hare, G. M. 2014. Differential HIF and NOS responses to acute anemia: defining organ-specific hemoglobin thresholds for tissue hypoxia. *Am J Physiol Regul Integr Comp Physiol*, 307, R13-25.
- Vandesompele, J., De Preter, K., Pattyn, F., Poppe, B., Van Roy, N., De Paepe, A. & Speleman, F. 2002. Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biol*, 3, RESEARCH0034.

- Viboolvorakul, S. & Patumraj, S. 2014. Exercise training could improve age-related changes in cerebral blood flow and capillary vascularity through the upregulation of VEGF and eNOS. *Biomed Res Int*, 2014, 230791.
- Vigano, A., Ripamonti, M., De Palma, S., Capitanio, D., Vasso, M., Wait, R., Lundby, C., Cerretelli, P. & Gelfi, C. 2008. Proteins modulation in human skeletal muscle in the early phase of adaptation to hypobaric hypoxia. *Proteomics*, 8, 4668-79.
- Vogt, M., Puntchart, A., Geiser, J., Zuleger, C., Billeter, R. & Hoppeler, H. 2001. Molecular adaptations in human skeletal muscle to endurance training under simulated hypoxic conditions. *J Appl Physiol*, 91, 173-82.
- Wang, G. L. & Semenza, G. L. 1995. Purification and characterization of hypoxia-inducible factor 1. *J Biol Chem*, 270, 1230-7.
- Wenger, R. H. & Gassmann, M. 1997. Oxygen(es) and the hypoxia-inducible factor-1. *Biol Chem*, 378, 609-16.
- Zoll, J., Ponsot, E., Dufour, S., Doutreleau, S., Ventura-Clapier, R., Vogt, M., Hoppeler, H., Richard, R. & Fluck, M. 2006. Exercise training in normobaric hypoxia in endurance runners. III. Muscular adjustments of selected gene transcripts. *J Appl Physiol*, 100, 1258-66.

Table 1. PCR Primers sequences.

Gene	Forward Primer	Reverse Primer
HIF-1 $\alpha$	GCCCCAGATTCAGGATCAGA	TGGGACTATTAGGCTCAGGTGAAC
PHD-2	AGGTGAAGCCAGCCTAT	CCTGATGCTAGCTGATACTTG
VEGF	TTTCTGCTGTCTTGGGTGCATTGG	ACCACTTCGTGATGATTCTGCCCT
Mb	GCCACCAAGCACAAAGATC	GGCATCAGCACCAAAGT
COX-4_1	GAGAGCTTTGCTGAGATGAA	CCGTACACATAGTGCTTCTG
COX-4_2	CCTTCTGCACAGAACTCAAC	CGGTACAAGGCCACCTTT
PGC-1 $\alpha$	GGGATGATGGAGACAGCTATGG	CTCTTGGTGGAAGCAGGGTC
TFAM	AGCGTTGGAGGGAAGTTCCTGATT	TTCTTTATATACCTGCCACTCCGCCC
eNOS	CAGTTACCAGCTAGCCAAAGT	CTCATTCTCCAGGTGCTTCAT
nNOS	CAGAACTCACACAAGGTCTATC	GTTGACCGACTGGATTTAGG
PDK-1	TGCCCATATCACGTCTTTAC	GTCTGTTGACAGAGCCTTAAT
GLUT-1	CCTGCAGTTTGGCTACAACA	GTGGACCCATGTCTGGTTG
PFK	ATTTGACGAAGCCCTGAAG	GTGCGAACCCTCTTAGATAC
RPL19	CGCTGTGGCAAGAAGAAGGTC	GGAATGGACCGTCACAGGC
B2M	ATGAGTATGCCTGCCGTGTGA	GGCATCTTCAAACCTCCATG

Table 2. Citrate synthase protein activity.

	LHTLH	LHTL	LLTL
Pre-	1.00 $\pm$ 0.00	1.00 $\pm$ 0.00	1.00 $\pm$ 0.00
Post-1	1.19 $\pm$ 0.27*, $\dagger$	0.82 $\pm$ 0.25	1.22 $\pm$ 0.25*, $\dagger$
Post-2	1.20 $\pm$ 0.31 $\dagger$	0.79 $\pm$ 0.32*	1.35 $\pm$ 0.22***, $\dagger$

\*  $P < 0.05$  \*\*\*  $P < 0.001$  significantly different from Pre-;  $P < 0.05$   $\dagger$  significantly different from LHTL.

## Figures Legend

Figure 1. Relative changes in selected mRNA expression markers from before (Pre-) to immediately (Post-1) and 3 weeks (Post-2) after the intervention. The intervention consisted in 14-days of passive normobaric hypoxic exposure combined with (a) repeated maximal-intensity hypoxic exercise in hypoxia (LHTLH) or (b) normoxia (LHTL). (c) A control group followed a ‘live low-train low’ (LLTL) protocol. Black and grey bars represent Post-1 and Post-2 values of mRNA concentrations in *vastus lateralis* muscle, respectively. These values were normalized to baseline values (Pre-), which were set to  $1.00 \pm 0.00$  (dashed line). Values are means  $\pm$  SD. \*  $P < 0.05$ , \*\*  $P < 0.01$  vs. Pre-intervention; #  $P < 0.05$  vs. LLTL and †  $P < 0.05$  vs. LHTL. HIF-1 $\alpha$ , hypoxia inducible factor-1 $\alpha$ ; PHD-2, prolyl hydroxylase domain protein 2; VEGF, vascular endothelial growth factor; Mb, myoglobin; PGC1- $\alpha$ , proliferator-activated receptor gamma coactivator-1 $\alpha$ ; TFAM, mitochondrial transcription factor A; COX-4\_1, cytochrome oxidase 4 isoform 1; COX-4\_2, cytochrome oxidase 4 isoform 2; PDK-1, pyruvate dehydrogenase kinase 1; GLUT-1, glucose transporter 1; PFK, phosphofructokinase; eNOS, endothelial nitric oxide synthase; nNOS, neuronal nitric oxide synthase.

Figure 2. Relative protein expression of selected markers from baseline (Pre-) to the end of the intervention (Post-1) and after 3 weeks (Post-2). The intervention consisted in 14-days of passive normobaric hypoxic exposure combined with (a) repeated maximal-intensity hypoxic exercise in hypoxia (LHTLH) or (b) normoxia (LHTL). (c) A control group followed a ‘live low-train low’ (LLTL) protocol. Black and grey bars represent Post-1 and Post-2 values of protein concentrations in *vastus lateralis* muscle, respectively, and were normalized to baseline values (Pre-), which were set to  $1.00 \pm 0.00$  (dashed line). Values are means  $\pm$  SD. \*  $P < 0.05$ , \*\*  $P < 0.01$ , \*\*\*  $P < 0.001$ , vs. Pre- and #  $P < 0.05$ , ##  $P < 0.01$  vs. LLTL. CS,

citrate synthase; MCT-1, monocarboxylate transporter 1; MCT-4, monocarboxylate transporter 4; CA-3, carbonic anhydrase III; GLUT-4, glucose transporter 4; PFK, phosphofructokinase; AMPK, AMP-activated protein kinase.

Figure 3. Representative Western blots showing the content of selected proteins from baseline (Pre-) to the end of the intervention (Post-1) and after 3 weeks (Post-2). The intervention consisted in 14-days of passive normobaric hypoxic exposure combined with (a) repeated maximal-intensity hypoxic exercise in hypoxia (LHTLH) or (b) normoxia (LHTL). (c) A control group followed a ‘live low-train low’ (LLTL) protocol. CS, citrate synthase; MCT-1, monocarboxylate transporter 1; MCT-4, monocarboxylate transporter 4; CA-3, carbonic anhydrase III; GLUT-4, glucose transporter 4; PFK, phosphofructokinase; AMPK, AMP-activated protein kinase.



